

International Journal of Modern Physics D
 © World Scientific Publishing Company

Magnetars and White Dwarf Pulsars

Ronaldo V. Lobato[†] and Manuel Malheiro^{*}

*Departamento de Física, Instituto Tecnológico de Aeronáutica,
 São José dos Campos, 12245030, São Paulo, Brazil
 E-mail: [†]rvlobato@ita.br; ^{*}malheiro@ita.br
<http://www.ita.br/>*

Jaziel G. Coelho

*Divisão de Astrofísica, Instituto Nacional de Pesquisas Espaciais - DAS/INPE/MCTI,
 São José dos Campos, 12227-010, São Paulo, Brazil
 E-mail: jaziel.coelho@inpe.br
<http://www.das.inpe.br/>*

Received Day Month Year

Revised Day Month Year

The Anomalous X-ray Pulsars (AXPs) and Soft Gamma-ray Repeaters (SGRs) are a class of pulsars understood as neutron stars (NSs) with super strong surface magnetic fields, namely $B \gtrsim 10^{14}$ G, and for that reason are known as Magnetars. However, in the last years some SGRs/AXPs with low surface magnetic fields $B \sim (10^{12} - 10^{13})$ G have been detected, challenging the Magnetar description. Moreover, some fast and very magnetic white dwarfs (WDs) have also been observed, and at least one showed X-Ray energy emission as an ordinary pulsar. Following this fact, an alternative model based on white dwarfs pulsars has been proposed to explain this special class of pulsars. In this model, AXPs and SGRs as dense and magnetized white dwarfs can have surface magnetic field $B \sim 10^7 - 10^{10}$ G and rotate very fast with frequencies $\Omega \sim 1$ rad/s, consistent with the observed rotation periods $P \sim (2 - 12)$ s.

Keywords: magnetic fields; magnetic white dwarfs; pulsars; SGRs/AXPs

PACS numbers: 97.60.Gb, 97.60.Jd, 04.40.Dg, 98.70.Qy

1. Introduction

The Soft Gamma-ray Repeaters and Anomalous X-ray Pulsars are a special class of pulsars, they are understood in the framework of strongly magnetized neutron star,^{1,2} but there are alternative scenarios, in particular the white dwarf (WD) pulsar model developed by Malheiro, Rueda and Ruffini.³ Recently, four over a total of about 23 SGRs/AXPs presented radio-pulsed emission. These radio sources showed several properties that make them different from the others SGRs/AXPs.

As pointed by Coelho and Malheiro⁴ the large steady X-ray luminosity seen for some no-radio SGRs/AXPs, can be explained as coming from a large spin-down energy lost of a massive white dwarf with a much large magnetic dipole

moment of $10^{34} \leq \mu \leq 10^{36}$ emu consistent with the range observed for isolated and very magnetic WDs (see Coelho & Malheiro 2012⁵ for discussions), indicating a different nature between these sources and the radio SGRs/AXPs. These radio SGRs/AXPs have large magnetic field and seem to be very similar to the high-B pulsars recently founded.⁶ However, even if the radio SGRs/AXPs are strong magnetized neutron stars, they are not magnetars in the sense that their steady luminosity is not originated by the magnetic energy, but from the rotational energy as rotation powered pulsars.

2. Canonical spin-powered pulsar model

Rotation powered pulsars (RPPs) are detected by its radio emission and their luminosity L_X is considered as coming from rotation.

In this model we consider a dipole moment μ with an orientation α in the rotation axis that rotates with an angular frequency Ω . On the surface of the star the magnetic field at the equator is $B_s \sim \mu/R^3$ and at the poles $B_p = 2\mu/R^3$, where R is the star radius. The rotational energy is given by,⁷

$$E_{\text{rot}} = \frac{1}{2} I \Omega^2, \quad (1)$$

where I is the moment of inertia. Thus, the loss of rotational energy of the pulsar is:

$$\dot{E}_{\text{rot}} = I \Omega \dot{\Omega} + \frac{1}{2} \dot{I} \Omega^2 \approx I \Omega \dot{\Omega}. \quad (2)$$

Such configuration has a time-varying magnetic dipole moment, in an infinity referential, and radiates energy with a rate

$$\dot{E}_{\text{dip}} = \frac{2\mu^2 \Omega^4 \sin^2 \alpha}{3c^3}, \quad (3)$$

where c is the speed of light. Combining (2) and (3) we obtain the surface magnetic field at the equator as:⁸

$$B_s = B_p/2 = \left(\frac{3c^3 I}{8\pi^2 R^6} P \dot{P} \right)^{1/2}, \quad (4)$$

with period $P = 2\pi/\Omega$, and $\dot{P} = dP/dt$. We see that the value of the magnetic field is inferred from $\sqrt{P\dot{P}}$ that are measures obtained by astronomical observations, and two parameters, the moment of inertia I and the star radius R that depend how the mass-energy density is distributed inside the star, are obtained solving the Tolman-Oppenheimer-Volkoff equation for a specific equation of state chosen, and of course, are quite different for Neutron Stars or White Dwarfs.

3. The magnetar model

In the magnetar model, the SGRs/AXPs are considered as neutron stars with large magnetic fields, higher than the value of critical field from the quantum electrodynamics (QED) $B_q \equiv m^2 c^3 / \hbar e = 4.4 \times 10^{13}$ G (m is the electron's mass, c the speed of light and $\hbar$ the reduced Planck's constant and e the charge) and 100-1000 times than the magnetic field of normal pulsars observed. The observed X-ray luminosity L_X is determined by complex structures in its magnetosphere (see^{1,2}), considering that luminosity is powered by the energy of magnetic field.

The recent discovery of radio-pulsed emission in four of this class of sources, where the spin-down rotational energy lost $\dot{E}_{\text{rot}}$ is larger than the X-ray luminosity L_X as we see in the Figure (1) opens the question of the nature of these radio sources in comparison to the other SGRs/AXPs (we have also recently studied this point in Refs.,⁹⁻¹¹ where the models of radio emission were investigated).

3.1. Difficulties with the magnetar model

According the magnetar model the high X-ray luminosity from this sources is powered by the energy stored in their strong magnetic fields.^{12,13} Although the interpretation as neutron stars is the one most successfully, here are some aspects about SGRs/AXPs not very well understood in this interpretation:

- RPPs are assumed born with a short period ~ 1 ms resulting in large magnetic fields by dynamo process,¹⁴ the SGRs/AXPs presents long period, such that generation of high magnetic fields by this mechanisms is not well understood.
- the large spin-down rates in comparison with the typical pulsar implies, for example, that the source SGR 0418+5729 has a characteristic age $\tau = P/2\dot{P} = 2.4 \times 10^7$ years. Thus, it is difficult to understand, how being much older than ordinary pulsars, it still fits in the magnetar model where the sources are seen as young pulsars.

In fact, some SGRs/AXPs with low magnetic fields and smaller spin-down rates were observed.¹⁵ In this observation we highlight SGR 0418+5729, with period $P = 9.08$ s, spin-down $\dot{P} < 6.0 \times 10^{-15}$ s/s and X-ray luminosity $L_X < 6.2 \times 10^{31}$ erg/s, considering this source a neutron star, the loss of energy associated with injection of rotational energy in the pulsars magnetosphere does not explain the luminosity of this star, i.e., $\dot{E}_{\text{rot}}^{\text{NS}} < L_X$. Thus, excludes the possibility to be a RPP, where $\dot{E}_{\text{rot}}^{\text{NS}} > L_X$. Recent observations of Fermi satellite¹⁶⁻¹⁸ does not find evidences of γ radiation. Furthermore, were detected 4 SGRs/AXPs that emit in wavelength of radio, while for the others not. These observations carried some uncertainties about the nature of SGRs/AXPs and opens the possibility of consider a different nature for these sources, as white dwarf pulsars as proposed by Malheiro, Rueda and Ruffini.³

4. SGRs/AXPs as white dwarf pulsars

Rapidly, massive white dwarfs and with observed high magnetic field $10^6 - 10^9$ G were detected in the recent years¹⁹ and share some similarities with SGRs/AXPs. After these facts, Malheiro and collaborators³ following two works from Morini²⁰ and Paczynski²¹ developed an alternative model explaining SGRs/AXPs as white dwarfs. As was investigated by Usov in 1989,²² the process of release energy in a massive, magnetic white dwarf can be explained in terms of a canonical spin-powered pulsars model, because in several aspects they are similar.

For example, if we consider a star with $M = 1.4M_\odot$ and $R = 10^6$ cm (a neutron star), from (4) the magnetic field at poles is:

$$B_p^{\text{NS}} = 3.2 \times 10^{19} (P\dot{P})^{1/2} \text{G}, \quad (5)$$

In the case of a white dwarf with $M = 1.4M_\odot$ and $R = 3 \times 10^8$ cm, there is a new scale for the magnetic field at poles:

$$B_p^{\text{WD}} = 4.21 \times 10^{14} (P\dot{P})^{1/2} \text{G}, \quad (6)$$

and to the dipole moment μ , responsible by the dipole radiation emitted, which is a factor 10^3 times larger than for neutron stars. That is the factor observed between the X-Ray luminosity of SGRs/AXPs as white dwarfs and slowly pulsars like the XDINS,²³ and high magnetic pulsars:²⁴ basically all these sources present the same period $P \sim 1 - 10$ s that SGRs/AXPs. When SGRs/AXPs are white dwarfs there are new values for the mass density, moment of inertia, dipole moment and rotation energy. It explains a large class of problems related in considering SGRs/AXPs as neutron stars. Furthermore, we cannot ignore the recent astronomical observations of old SGRs (characteristic age $\sim (10^6 - 10^7)$ Myr) with low surface magnetic field, as well as the four SGRs/AXPs showing radio emission recently discussed by Coelho and Malheiro.^{25,26} In the table (1) we summarized the properties among SGR 0418+5729, Swift J1822.3-1606²⁷ (both SGRs with low-B), AE Aquarii²⁸ (the first white dwarf pulsar), RXJ 0648.0-4418 (rapid WD that emit in X-ray), and the candidate EUVE J0317-855.²⁹

Table 1. Comparison between the observational and deduce proprieties in the dipole model, where the SGRs/AXPs were consider as white dwarfs: the quantities P , $\dot{P}$ and L_X were obtained in.³⁰

Stars	P (s)	$\dot{P}$ (10^{-14} s/s)	τ (10^6 years)	L_X (erg/s)	B_e^{WD} (G)	μ_p^{WD} (emu)
SGR 0418	9.08	< 0.6	24	$\sim 6.2 \times 10^{31}$	$< 9.83 \times 10^7$	2.65×10^{33}
Swift J1822	8.44	8.3	1.6	$\sim 4.2 \times 10^{32}$	3.52×10^8	0.95×10^{34}
AE Aquarii	33.08	5.64	9.3	$\sim 10^{31}$	$\sim 5 \times 10^7$	$\sim 1.35 \times 10^{33}$
RX J0648	13.2	0.6	0.23	$\sim 10^{32}$	0.1×10^9	3.48×10^{34}
EUVE J0317	725	-	-	-	$\sim 4.5 \times 10^8$	1.22×10^{34}

In the description of SGRs/AXPs as white dwarfs some problems inherent to magnetar models are solved as consequence of the new scales:

- the existence of stable white dwarfs could explain the band of period $2 \lesssim P \lesssim 12$ s observed in SGRs/AXPs. In particular the fact that we do not observe SGRs/AXPs with $P < 2$ s is due the white dwarf structure. In³¹ self-consistent computations shown that the minimum period to white dwarfs are $\sim 0.3, 0.5, 0.7$ and 2.2 s to star made of ^4He , ^{12}C , ^{16}O , ^{56}Fe respectively.
- the energy balance is solved, since the X-luminosity observed is less than the rotational energy of a white dwarf, $L_X < \dot{E}_{\text{rot}}$, because $\dot{E}_{\text{rot}}$ is 10^5 bigger than the value to a neutron star. SGRs/AXPs powered by rotational energy are completely analogous to RPPs of neutron star.
- largest luminosity $L_X \sim 10^{35}\text{erg/s}$ to slowly pulsar ($\Omega \sim 1\text{Hz}$) is understood as consequence of the white dwarfs' radius that produces a large dipole moment. Some works shown the possibility of pulsed radiation in white dwarfs. SGRs/AXPs like white dwarfs have large magnetic fields and rotate very fast, near the Kepler frequency, and they could produce large potential difference on its surface able to produce X and γ radiation.³² Bursts in this sources, that is common, would be a consequence of their angular velocity and change in the moment of inertia. In this situation there are blunt changes in gravitational energy.³
- SGRs/AXPs has a low population in contrast with more than 2000 RPPs founded, this is justified by the motivation that white dwarfs with high magnetic field an rapidly are rarely formed.³³ Furthermore following the observations, white dwarfs with high magnetic fields are only 10% of magnetic stars.

4.1. Efficiency

SGRs/AXPs when understood as neutron stars typically have a larger luminosity in X-ray that can not be explained by their spin-down luminosity, i.e., $L_X \gg \dot{E}_{\text{rot}}$. However, in four of them that is not true and they present radio emission. These radio SGRs/AXPs showed several properties different from the others SGRs/AXPs. As was discussed, in this four sources the X-ray efficiency $\eta_X = L_X/\dot{E}_{\text{rot}}$ seems to be small comparing with SGRs/AXPs as white dwarfs.^{3,26} However, when these four sources are understood as neutron stars pulsars the efficiency is $\eta_X \sim (0.2-0.1)$, much larger than the values of ordinary pulsars where $\eta_X \sim (10^{-3} \text{ to } 10^{-4})$, but still smaller than one: for this reason Coelho and Malheiro²⁶ suggested that these four star are very efficient neutron stars RPPs in contrast to the others that are white dwarfs RPPs. As we can see in the figure (1) this four sources presents a loss of rotational energy larger than the X-ray luminosity. They are very similar to the high- B pulsars recently found^{34,35} which have their luminosity powered by rotational energy.




Fig. 1. Four SGRs/AXPs in comparison with all no-radio SGRs/AXPs as neutron stars. These radio sources have large magnetic fields and are very similar to high- B pulsars. Even this sources are strong magnetized neutron stars, they are not magnetar, because their steady luminosity can be explained by loss of rotational energy.

4.2. *Difficulties with the white dwarf pulsar model*

Although SGRs/AXPs as white dwarfs pulsars are well described, there are some difficulties aspects in the phenomenology:

- Some SGRs/AXPs are associate with supernova. Usually, this association is an evidence of SGRs/AXPs to be neutron stars; and also some are associate with new born stars' clusters.
- AXPs show up variation in its luminosity and some transients seems to contradict the rotation powered hypotheses.

SGRs/AXPs as white dwarfs pulsars in comparison with magnetar have larger radius. However is difficult to estimate the radius due to the distance, since the majority of SGRs/AXPs are in $d \gtrsim 2$ kpc. The knowledge about their nature has an uncertainty until we have better measures for the radii of these sources.

5. Conclusions

Some SGRs/AXPs within the description of white dwarfs pulsars can be well understood. In the white dwarfs pulsars scenario we solve the majority of problems concerning the magnetar model, in particular the ultra high magnetic fields. Super-massive, fast and highly magnetized white dwarfs have already been found and, this number is increasing every year, opening a new research area. The future will solve the correct nature of SGRs/AXPs: Magnetars or White Dwarf pulsars.

Acknowledgments

This work was financially supported by CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) and FAPESP thematic project 2015/26258-4. J.G.C. acknowledges the support of FAPESP (2013/15088-0 and 2013/26258-4). Acknowledgements also to organizing committee for the hospitality in the III Amazonian Symposium on Physics and V NRHEP Network Meeting: Celebrating 100 years of General Relativity.

References

1. R. C. Duncan and C. Thompson, Formation of very strongly magnetized neutron stars - implications for gamma-ray bursts, *The Astrophysical Journal Letters* **392**, L9 (6 1992).
2. C. Thompson and R. C. Duncan, The soft gamma repeaters as very strongly magnetized neutron stars - i. radiative mechanism for outbursts, *Monthly Notices of the Royal Astronomical Society* **275**, 255 (7 1995).
3. M. Malheiro, J. A. Rueda and R. Ruffini, {SGRs} and {axps} as rotation-powered massive white dwarfs, *Publications of the Astronomical Society of Japan* **64**, p. 56 (6 2012).
4. J. G. Coelho and M. Malheiro, Magnetic dipole moment of sgrs and axps described as massive and magnetic white dwarfs, *Publications of the Astronomical Society of Japan* **66**, 14 (2 2014).
5. J. G. Coelho and M. Malheiro, Similarities of sgrs with low magnetic field and white dwarf pulsars, *International Journal of Modern Physics: Conference Series* **18**, 96 (1 2012).
6. N. Rea, J. A. Pons, D. F. Torres and R. Turolla, The fundamental plane for radio magnetars, *The Astrophysical Journal Letters* **748**, p. L12 (3 2012).
7. S. L. Shapiro and S. A. Teukolsky, *Black Holes, White Dwarfs and Neutron Stars: The Physics of Compact Objects* (John Wiley & Sons, 11 2008).
8. A. Ferrari and R. Ruffini, Theoretical implications of the second time derivative of the period of the pulsar np 0532, *The Astrophysical Journal Letters* **158**, p. L71 (11 1969).
9. J. G. Coelho and M. Malheiro, Sgrs and axps as white dwarf pulsars (AIP Publishing, 3 2013).
10. R. V. Lobato, J. Coelho and M. Malheiro, Particle acceleration and radio emission for sgrs/axps as white dwarf pulsars, *Journal of Physics: Conference Series* **630**, p. 012015 (7 2015).

8 *R. V. Lobato, M. Malheiro, J. G. Coelho*

11. R. V. Lobato, J. G. Coelho and M. Malheiro, Radio pulsar death lines to sgrs/axps and white dwarfs pulsars, in *THE SECOND ICRANET CÉSAR LATTES MEETING: Supernovae, Neutron Stars and Black Holes*, (AIP Publishing, 2015).
12. S. Mereghetti, Pulsars and magnetars, *arXiv:1304.4825 [astro-ph]* (4 2013).
13. G. Giovannetti, D. Puggioni, J. M. Rondinelli and M. Capone, Electronic correlations and screening effects in the hund's polar metal sreumo\$_{2}\$so\$_{6}\$, *arXiv:1507.02924 [astro-ph]* (7 2015).
14. H. Tong and R.-X. Xu, Magnetars: fact or fiction?, *International Journal of Modern Physics E* **20**, 15 (12 2011).
15. N. Rea, P. Esposito, R. Turolla, G. L. Israel, S. Zane, L. Stella, S. Mereghetti, A. Tiengo, D. G. E. G and C. Kouveliotou, A low-magnetic-field soft gamma repeater, *Science* **330**, 944 (11 2010).
16. A. A. Abdo, M. Ackermann, M. Ajello, A. Allafort, L. Baldini, J. Ballet, G. Barbiellini, M. G. Baring, D. Bastieri, R. Bellazzini, R. D. Blandford, E. D. Bloom, E. Bonamente, A. W. Borgland, A. Bouvier, J. Bregeon, M. Brigida, P. Bruel, T. H. Burnett, G. A. Caliendo, R. A. Cameron, P. A. Caraveo, C. Cecchi, O. Celik, S. Chaty, A. Chekhtman, C. C. Cheung, J. Chiang, S. Ciprini, R. Claus, J. Conrad, P. R. d. Hartog, C. D. Dermer, A. d. Angelis, F. d. Palma, R. Dib, M. Dormody, E. do Couto e Silva, P. S. Drell, R. Dubois, D. Dumora, T. Enoto, C. Favuzzi, M. Frailis, P. Fusco, F. Gargano, N. Gehrels, N. Giglietto, P. Giommi, F. Giordano, M. Giroletti, T. Glanzman, G. Godfrey, I. A. Grenier, M.-H. Grondin, S. Guiriec, D. Hadasch, Y. Hanabata, A. K. Harding, E. Hays, G. L. Israel, G. Jóhannesson, A. S. Johnson, V. M. Kaspi, H. Katagiri, J. Kataoka, J. Knödseder, M. Kuss, J. Lande, S.-H. Lee, M. Lemoine-Goumard, F. Longo, F. Loparco, M. N. Lovellette, P. Lubrano, A. Makeev, M. Marelli, M. N. Mazziotta, J. E. McEnery, J. Mehault, P. F. Michelson, T. Mizuno, A. A. Moiseev, C. Monte, M. E. Monzani, A. Morselli, I. V. Moskalenko, S. Murgia, M. Naumann-Godo, P. L. Nolan, E. Nuss, T. Ohsugi, A. Okumura, N. Omodei, E. Orlando, J. F. Ormes, M. Ozaki, D. Paneque, D. Parent, M. Pepe, M. Pesce-Rollins, F. Piron, T. A. Porter, S. Rainò, R. Rando, M. Razzano, N. Rea, A. Reimer, O. Reimer, T. Reposeur, S. Ritz, H. F.-W. Sadrozinski, P. M. S. Parkinson, C. Sgrò, E. J. Siskind, D. A. Smith, G. Spandre, P. Spinelli, M. S. Strickman, H. Takahashi, T. Tanaka, J. B. Thayer, D. J. Thompson, L. Tibaldo, D. F. Torres, G. Tosti, A. Tramacere, E. Troja, Y. Uchiyama, T. L. Usher, J. Vandenbroucke, V. Vasileiou, G. Vianello, V. Vitale, A. P. Waite, B. L. Winer, K. S. Wood, Z. Yang and M. Ziegler, Search for gamma-ray emission from magnetars with the fermi large area telescope, *The Astrophysical Journal Letters* **725**, p. L73 (12 2010).
17. H. Tong, L. M. Song and R. X. Xu, Non-detection in a fermi/lat observation of axp 4u 0142+61: Magnetars?, *The Astrophysical Journal Letters* **725**, p. L196 (12 2010).
18. H. Tong, L. M. Song and R. X. Xu, Anomalous x-ray pulsars and soft gamma-ray repeaters in the outer gap model: Confronting fermi observations, *The Astrophysical Journal* **738**, p. 31 (9 2011).
19. B. G. Castanheira, S. O. Kepler, S. J. Kleinman, A. Nitta and L. Fraga, Discovery of five new massive pulsating white dwarf stars, *Monthly Notices of the Royal Astronomical Society* **430**, 50 (3 2013).
20. M. Morini, N. R. Robba, A. Smith and M. van der Klis, Exosat observations of the supernova remnant g109.1-1.0 and the x-ray pulsar 1e 2259+586, *The Astrophysical Journal* **333**, 777 (10 1988).
21. B. Paczynski, X-ray pulsar 1e 2259 + 586 - a merged white dwarf with a 7 second rotation period?, *The Astrophysical Journal Letters* **365**, L9 (12 1990).
22. V. Usov, Gamma-radiation generation by rotating magnetic white dwarfs, *Pisma v*

- Astronomicheskii Zhurnal* **14**, p. 606609 (1988).
23. D. L. Kaplan and M. H. v. Kerkwijk, Constraining the spin-down of the nearby isolated neutron star rx j2143.0+0654, *The Astrophysical Journal Letters* **692**, p. L62 (2 2009).
 24. F. P. Gavril, M. E. Gonzalez, E. V. Gotthelf, V. M. Kaspi, M. A. Livingstone and P. M. Woods, Magnetar-like emission from the young pulsar in kes 75, *Science* **319**, 1802 (3 2008).
 25. J. G. Coelho and M. Malheiro, A different nature between the radio axps in comparison to the others sgrs/axps, *arXiv:1307.8158 [astro-ph, physics:nucl-th]* (7 2013).
 26. J. G. Coelho and M. Malheiro, Do sgrs/axps and radio axps have the same nature?, *arXiv:1303.0863 [astro-ph]* (3 2013).
 27. N. Rea, G. L. Israel, J. A. Pons, R. Turolla, D. Viganò, S. Zane, P. Esposito, R. Perna, A. Papitto, G. Terreran, A. Tiengo, D. Salvetti, J. M. Girart, A. Palau, A. Possenti, M. Burgay, E. G. G. A. Caliendo, C. Kouveliotou, D. G. R. P. Mignani, E. Ratti and L. Stella, The outburst decay of the low magnetic field magnetar sgr 0418+5729, *The Astrophysical Journal* **770**, p. 65 (6 2013).
 28. Y. Terada, T. Hayashi, M. Ishida, K. Mukai, T. Dotani, S. Okada, R. Nakamura, S. Naik, A. Bamba and K. Makishima, Suzaku discovery of hard x-ray pulsations from a rotating magnetized white dwarf, aequarii, *Publications of the Astronomical Society of Japan* **60**, 387 (4 2008).
 29. L. Ferrario, S. Vennes, D. T. Wickramasinghe, J. A. Bailey and D. J. Christian, Euve j0317-855 a rapidly rotating, high-field magnetic white dwarf, *Monthly Notices of the Royal Astronomical Society* **292**, p. 205 (11 1997).
 30. S. A. Olausen and V. M. Kaspi, The mcgill magnetar catalog, *The Astrophysical Journal Supplement Series* **212**, p. 6 (5 2014).
 31. K. Boshkayev, J. A. Rueda, R. Ruffini and I. Siutsou, On general relativistic uniformly rotating white dwarfs, *The Astrophysical Journal* **762**, p. 117 (1 2013).
 32. K. Kashiyama, K. Ioka and P. Mészáros, Cosmological fast radio bursts from binary white dwarf mergers, *The Astrophysical Journal Letters* **776**, p. L39 (10 2013).
 33. M. Ilkov and N. Soker, Type ia supernovae from very long delayed explosion of core-white dwarf merger, *Monthly Notices of the Royal Astronomical Society* **419**, 1695 (1 2012).
 34. S. A. Olausen, V. M. Kaspi, A. G. Lyne and M. Kramer, Xmm-newton x-ray observation of the high-magnetic-field radio pulsar psr j1734-3333, *The Astrophysical Journal* **725**, p. 14 (12 2010).
 35. S. A. Olausen, W. W. Zhu, J. K. Vogel, V. M. Kaspi, A. G. Lyne, C. M. Espinoza, B. W. Stappers, R. N. Manchester and M. A. McLaughlin, X-ray observations of high-b radio pulsars, *The Astrophysical Journal* **764**, p. 1 (2 2013).